Winter Run Chinook Salmon Draft Preliminary Example Biological Goals, Objectives, and Stressors Logic Chain for the BDCP February 27th, 2012

Global Goal

Removal of the Sacramento River winter-run Chinook salmon ESU from the Federal List of Endangered and Threatened Wildlife (NMFS 2009). According to the NMFS draft recovery plan (2009), recovery and long-term sustainability requires:

- 1) Adequate protection for replacement of losses due to natural mortality (disease and stochastic events);
- 2) Sufficient genetic robustness to avoid inbreeding depression and allow for adaptation
- 3) Sufficient habitat (type, amount, and quality) for long-term population maintenance, and
- 4) Elimination or control of threats.

Global Objectives:

There are two components of Global Objectives that are relevant to the BDCP program. The first pertains to Recovery Plan goals based on Viable Salmonid Population (VSP) criteria, and further refined for the Central Valley in Lindley et al. (2007). The second component relates to Critical Habitat (as designated for Winter-run Chinook June 16, 1993; 58 FR 33212)

Global VSP Objectives include:

- VSP1. Increase abundance
- VSP2. Increase spatial distribution
 - a. Secure all extent populations (all populations are important because there are so many "missing" populations in the Central Valley)
 - b. Recover populations in each diversity group (only one diversity group for Winter-run)
- VSP3. Protect and increase life history and genetic diversity
- VSP4. Increase productivity (population growth rate = births-deaths)

Viable populations should demonstrate a combination of population growth rate and abundance that produces an acceptable probability of population persistence (NMFS Draft Recovery Plan).

Global Critical Habitat Objectives (from primary constituent elements)

- CH1. Provide access to spawning areas on the Upper Sacramento River, including upstream passage of adults to spawning grounds
- CH2. Provide adequate quality and quantity of spawning gravels
- CH3. Provide for adequate river flows for successful spawning, incubations of eggs, fry development and emergence, and downstream transport of juveniles
- CH4. Provide water temperatures for successful spawning, egg incubation, and fry development
- CH5. Provide habitat areas and prey that are not contaminated
- CH6. Provide riparian (including floodplain) habitat for successful juvenile development and survival
- CH7. Provide adequate downstream migration corridors for successful emigration of juveniles

SPERT

| Global Goal | Global Objectives | BDCP Goal | BDCP Objective | Assumed Stressor | Stressor Reduction Target | Cons. Measures |
|--|---|--|---|--|---|--|
| Increase winter-run Chinook salmon abundance | Attainment of the winter- run Chinook salmon global abundance goal will occur by 2060 with | Improved juvenile survival (as a proxy for abundance) within the Plan Area, SF Bay, and the nearshore ocean. | Double current survival rate of juvenile emigrants from Sacramento River through Delta (35%) to 70%1. | Entrainment | Reduce entrainment of winter-run by at least 50% in all water year types | -Water Ops -Alternate migration routes |
| | achievement of 6-year geometric mean escapement levels of: 20,000 in the mainstem Sacramento River with | | | Predation | -Decrease mortality from predation in (specify locations & months) by%Increase quantity and quality of rearing habitat (including floodplain, channel margin, and riparian habitats) throughout the Delta | -Predator removal -Water Ops |
| | no year below 5,000; 3,000 in the Battle Creek watershed with no year below 700; and 800 in a third viable population with no year below 200. | | | Limited Rearing Habitat | Increase average size of juveniles (relative to current conditions) as they migrate through the Delta to% of their physiological maximum (corrected for temp) | -Yolo bypass -Suisun -Ammonia |
| | | | | North Delta Diversion Facilities | Maintain survival rates through the reach containing new north Delta diversions to no more than a 2% loss per screen, and no more than a 5% cumulative loss. | -Water Ops -Alternate migration routes -Predator removal |
| | | | | Limited Migration Flows | Maintain minimum Delta inflows in key migratory months | Water Ops |
| | | Increase migration/ spawning success | Eliminate human-induced passage delays | Migration barriers | Eliminate known human-caused passage impediments (chemical and physical) in the Plan Area | Fremont weir |
| | | of adult winter- run migrating through the Delta. | or illegal take of winter-run adults in the Delta. | Poaching | Eliminate winter-run poaching in the Delta | Funding for game wardens |

¹Estimate of 35% current survival is from Chinook Salmon Project Work Team. Survival varies by water year type, so survival would be measured

as a long-term average.

98JE/

| WINTER RUN – GLOBAL SPATIAL DISTRIBUTION GOAL LOGIC TREE | | | | | | | |
|---|---|---|---|------------------------------------|---|-----------------------------|--|
| Global | Global Objectives | BDCP Goal | BDCP | Assumed | Stressor Reduction Target | Cons. | |
| Goal | | | Objective | Stressor | | Measures | |
| Increase spatial distribution | Attainment of the winter-run Chinook salmon global spatial | Improved adult and juvenile migration success | Eliminate human-induced adult passage | Migration barriers | Eliminate known human-caused passage impediments (physical) within Plan Area | Fremont weir | |
| of winter- run Chinook salmon | distribution goal will occur by 2060 with restoration of two self- sustaining, independent | through the Delta to and from Sacramento River Basin (as a proxy | delays (barriers that necessitate median passage time > 36 hrs.) | Attraction flows | Provide Delta inflows >cfs betweendate &date | Operation s | |
| | populations in two watersheds of the Sacramento River drainage, and a third | for spatial distribution) | Create one alternate migratory path on the lower | Entrainment | Maintain Sac River winter-run entrainment at project pumps to < 2% of estimated smolt production in all water year types. | Water Ops | |
| | dependent population in the Sacramento River drainage. | | Sacramento River in > 40% of years. | Limited juvenile emigration routes | Ensure that at least % of juvenile winter-run emigrate through a non-mainstem channel route (e.g. flood bypass) in at least 40% of years. | -Yolo bypass -Fremont | |
| | | | | | | weir | |

| WINTER RUN – GLOBAL LIFE HISTORY AND GENETIC DIVERSITY GOAL LOGIC TREE | | | | | | | | |
|---|---|--|---|---------------------------------|---|--------------------------------------|--|--|
| Global Goal | Global Objectives | BDCP Goal | BDCP Objective | Assumed Stressor | Stressor Reduction Target | Cons. Measures | | |
| Conserve and restore life-history and genetic diversity of winter- run Chinook salmon | Protect and restore the full range of adult and juvenile life- history types migrating through the | Ensure that the project does not favor the survival of one life-history type | oes not artificial selection for winter-run life-history types resulting from | Hatchery Effects | Adopt hatchery practices that minimize adverse changes to life-history traits (e.g., size/age at smolting, age at maturity, migration timing) of wild winter-run Chinook. | Hatchery reform | | |
| | Delta. Attainment of the winter-run Chinook salmon global life history diversity goal will occur by 2060 with | over others. | | Hatchery Effects Entrainment | Alter hatchery practices to minimize adverse changes in life-history traits of hatchery winter-run Chinook. Ensure that entrainment does not favor the survival of one life-history type over other types (e.g., early or late migrating smolts or adults) | Hatchery reform Water Ops | | |
| | restoration of two self- sustaining, independent populations in two watersheds of the Sacramento River drainage, and a third | | O. R. R. R. | Flow Magnitude and Timing | To an equal degree across winter-run life-history types (e.g., size/age at smolting, age at maturity, migration timing) provide flows that support rearing and migration in all times and places where they occur. | Water Ops (including upstream) | | |
| | dependent population in the Sacramento River drainage. | | | Rearing Habitat | Winter-run Chinook will have access to inundated floodplains > 45 days in at least 1 of 3 years, | Yolo Bypass | | |